



SAFeway PHARMACY
2020 MARKET STREET
SAN FRANCISCO, CA 94114
(415)436-9032
#1507

Official Receipt - Please retain for tax or insurance

GOODGE, LESLIE (415)235-0698
400 DUBOCE AVE #419 05/72
SAN FRANCISCO, CA 94114

DR. LEE, SIMON [NW]
ST FRANCIS MEMORIAL
SAN FRANCISCO, CA 94109

Rx:6909549 Nov 30, 2006 Safety Cap: Yes
CIPROFLOXACIN 250MG TAB (RANB) Qty: 20 TAB
Generic for: CIPRO 250MG TAB

Ref: 062590534790987
ADVANCE PCS #610415 NDC: 63304-0709-01
Amount Due: \$5.00



REFILL YOUR PRESCRIPTIONS
@ SAFEWAY.COM

TAKE ONE TABLET BY MOUTH TWICE DAILY

Refills: 0

- 1-Drink plenty of fluids while on this medication
- 2-Do not take with antacids /iron/vitamins
- 3-Avoid large amnts of food or drinks with caffeine
- 4-if dizziness occurs, drive with caution.
- 5-Avoid prolonged exposure to sun. Use sunscreen
- 7-Promptly report tendon pain or swelling.



Rx 6909549

Safety Caps: Yes

Pat Allergies: IMIDAZOLES,



63304-0709-01 GOODGE, LESLIE

Labelsc.ro.02/1306

CIPROFLOXACIN 250MG TAB RANB

GENERIC NAME: CIPROFLOXACIN (sip-roe-FLOX-a-sin)

COMMON USES: This medicine is a fluoroquinolone antibiotic used to treat bacterial infections.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine comes with a patient information leaflet. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. This medicine may be taken on an empty stomach or with food. **DRINK A FULL GLASS OF WATER with each dose.** Drink several additional glasses of water daily, unless otherwise directed by your doctor. **DO NOT TAKE** any products containing magnesium, aluminum, or calcium within 6 hours before or 2 hours after taking this medicine. Some examples include quinapril, didanosine, vitamins/minerals, and antacids. Sulfafurazole, bismuth subsalicylate, iron, and zinc are also included. These products bind with ciprofloxacin preventing its full absorption. Avoid taking this medicine with milk or milk products (such as yogurt or calcium-enriched juice) by themselves. However, taking this medicine as part of a full meal that contains milk or milk products is permitted. Contact your doctor or pharmacist if you have any questions about how to properly take this medicine. **STORE THIS MEDICINE** at room temperature below 86 degrees F (30 degrees C), in a tightly-closed container, away from heat, moisture, and light. Do not store in the bathroom. Antibiotics work best when the amount of medicine in your body is kept at a constant level. Therefore, take this medicine at the same time each day. **TO CLEAR UP YOUR INFECTION COMPLETELY,** continue taking this medicine for the full course of treatment even if you begin to feel better in a few days. Do not miss any doses. **IF YOU MISS A DOSE OF THIS MEDICINE,** take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to this medicine or another fluoroquinolone antibiotic (such as levofloxacin, ofloxacin). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a fluoroquinolone, contact your doctor or pharmacist. **IF THIS MEDICINE CAUSES A SKIN RASH, HIVES, OR DIFFICULTY BREATHING,** check with your doctor before taking additional doses. **KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS** while you are using this medicine. **THIS MEDICINE MAY CAUSE DIZZINESS.** Taking this medicine alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. **THIS MEDICINE MAY CAUSE** increased sensitivity to the sun. Avoid exposure to the sun, sunlamps, or tanning booths until you know how you react to this medicine. Use a sunscreen or protective clothing if you must be outside for a prolonged period. **AVOID LARGE AMOUNTS OF CAFFEINE-CONTAINING** foods and beverages, such as coffee, tea, cocoa, cola drinks, and chocolate, while taking this medicine as the effects of the caffeine may be increased. This medicine may rarely cause pain, inflammation, or rupture of a tendon. If you experience any of these problems contact your doctor immediately. Rest and avoid exercise until you receive further instruction from your doctor. **BEFORE YOU BEGIN TAKING ANY NEW MEDICINE,** either prescription or over-the-counter, check with your doctor or pharmacist. Caution is advised when using this medicine in the elderly because they may be more sensitive to the side effects of the medicine (e.g., tendon problems, especially if they are also taking corticosteroid medicines such as prednisone). **IF YOU PLAN ON BECOMING PREGNANT,** discuss with your doctor the benefits and risks of using this medicine during pregnancy. **THIS MEDICINE IS EXCRETED IN BREAST MILK. DO NOT BREAST-FEED** while taking this medicine. **IF YOU HAVE DIABETES,** this medicine may affect your blood sugar. Check blood sugar levels closely and ask your doctor before adjusting the dose of your diabetes medicine.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include stomach upset, loss of appetite, diarrhea, nausea, headache, vision changes or dizziness during the first few days as your body adjusts to this medicine. If they continue or are bothersome, check with your doctor. **CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE** if you experience sunburn-like skin reaction (sun sensitivity) or severe or continuing diarrhea. **CONTACT YOUR DOCTOR IMMEDIATELY** if you experience seizures, mental or mood changes, easy bruising or bleeding, swelling of throat or tongue, persistent sore throat or fever, irregular heartbeat, chest pain, stomach pain, yellowing eyes and skin, dark urine, unusual change in the amount of urine or unusual fatigue. This medicine may rarely cause a severe intestinal condition (pseudomembranous colitis) due to a resistant bacteria. This condition may occur weeks after treatment has stopped. Do not use anti-diarrhea products or narcotic pain medicines because these products may make the effects worse. Therefore, **CONTACT YOUR DOCTOR IMMEDIATELY** if you develop tendon pain, joint pain or swelling, persistent diarrhea, abdominal or stomach pain or cramping, or blood or mucus in your stool. **USE OF THIS MEDICINE FOR PROLONGED OR REPEATED PERIODS** may result in oral thrush or a vaginal yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge or other new symptoms. **AN ALLERGIC REACTION** to this medicine is unlikely but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This information is generalized and is not intended as specific medical advice. If you have questions about the medicines you are taking or would like more information, check with your doctor, pharmacist, or nurse.