



SAFeway PHARMACY
2020 MARKET STREET
SAN FRANCISCO, CA 94114

#1507 (415)436-9032

Official Receipt - Please retain for tax or insurance

GOODGE, LESLIE (415)235-0698
400 DUBOCE AVE #419 05/72

SAN FRANCISCO, CA 94114

DR. LEE, SIMON [NW]

ST FRANCIS MEMORIAL

SAN FRANCISCO, CA 94109

Rx:6909551 Nov 30, 2006 Safety Cap: Yes

MALARONE 250/100 TAB (GLAX) Qty: 34 TAB

Ref: 06259055808831 NDC: 00173-0675-01 GH/SK

Amount Due: \$5.00



REFILL YOUR PRESCRIPTIONS

@ SAFEWAY.COM

TAKE ONE TABLET BY MOUTH ONE TIME DAILY

Qty Rem: 96 until Nov 29, 2007

- 1-Take with food or milk (or immediately after)
- 2-Finish all medication. Take on schedule.
- 3-Take this medicine at the same time every day.
- 4-May cause dizziness
- 5-Check with Dr. before taking any other medicine
- 6-May cause nausea/vomiting or other GI effects.
- 7-Report hives/itching/problems in breathing to Dr
- 8-Promptly report unusual symptoms/effects to Dr



Rx 6909551

Safety Caps: Yes

Pat Allergies: IMIDAZOLES,

CAUTION DO NOT TAKE OTHER MEDICINES
READ WITHOUT CHECKING WITH YOUR
PHYSICIAN, NURSE, PHARMACEUTICAL
SALES REPRESENTATIVE, OR PHARMACIST.



LabelSec.ro.02/1306

00173-0675-01 GOODGE, LESLIE

MALARONE 250/100 TAB GLAX

GENERIC NAME: ATOVAQUONE (a-TOE-va-kwone) and PROGUANIL (PRO-gwah-nill)

COMMON USES: This medicine is an antiprotozoal and antimalarial combination used to treat or prevent malaria.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. TAKE THIS MEDICINE with food or milk at the same time every day. If you vomit within 1 hour of taking your dose, take another dose. Inform your doctor that you will need more medicine to finish your treatment. STORE THIS MEDICINE at room temperature at 77 degrees F (25 degrees C) in a tightly closed container, away from heat and light. Brief storage between 59 and 86 degrees F (15 and 30 degrees C) is permitted. TO CLEAR UP YOUR INFECTION COMPLETELY, continue taking this medicine for the full course of treatment. Do not miss any doses. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible with food or milk. If it is almost time for your next dose, contact your doctor. DO NOT take 2 doses at once.

CAUTIONS: DO NOT TAKE THIS MEDICINE if you have had an allergic reaction to it or are allergic to any ingredient in this product. IF YOU EXPERIENCE difficulty breathing; tightness of chest; swelling of eyelids, face, or lips; or if you develop a rash or hives, tell your doctor immediately. Do not take any more doses of this medicine unless your doctor tells you to do so. This medicine helps to prevent malaria; however, it is still possible to contract the disease. Notify your doctor immediately that you may have been exposed to malaria if you develop flu-like symptoms including chills, fever, muscle aches, or headache. THIS MEDICINE MAY CAUSE dizziness. Do not drive, operate machinery, or do anything else that could be dangerous until you know how you react to this medicine. THIS MEDICINE MAY CAUSE vomiting. If it persists or becomes severe, contact your doctor. The medicine may not be absorbed if you are vomiting and therefore not be effective. Your doctor may suggest other medicines to treat the vomiting or treat your condition with a different medicine. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. Caution is advised when using this medicine in the elderly since they may be more sensitive to the effects of this medicine. FOR WOMEN: IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using this medicine during pregnancy. One of the ingredients in this medicine IS EXCRETED in breast milk. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may occur while taking this medicine include rash, nausea, diarrhea, headache, vomiting, fever, difficulty sleeping, stomach pain, mouth sores, or swelling or soreness of the mouth or tongue. If they continue or are bothersome, check with your doctor. CONTACT YOUR DOCTOR IMMEDIATELY if you experience loss of appetite, weight loss, unusual fatigue, yellowing of the eyes or skin, dark urine, dizziness, severe vomiting or diarrhea, fever, muscle or back pains. AN ALLERGIC REACTION TO THIS MEDICINE IS UNLIKELY, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This information is generalized and is not intended as specific medical advice. If you have questions about the medicines you are taking or would like more information, check with your doctor, pharmacist, or nurse.